


Unit

4



COORDINATE GEOMETRY

Unit Outcomes:

After completing this unit, you should be able to:

- ✚ apply the distance formula to find the distance between any two given points in the coordinate plane.
- ✚ formulate and apply the section formula to find a point that divides a given line segment in a given ratio.
- ✚ write different forms of equations of a line and understand related terms.
- ✚ describe parallel or perpendicular lines in terms of their slopes.

Main Contents

- 4.1 Distance between two points
- 4.2 Division of a line segment
- 4.3 Equation of a line
- 4.4 Parallel and perpendicular lines

Key Terms

Summary

Review Exercises

INTRODUCTION

IN **UNIT 3**, YOU HAVE SEEN AN IMPORTANT CONNECTION BETWEEN ALGEBRA AND GEOMETRY. THE GREAT DISCOVERIES OF THE 17TH CENTURY MATHEMATICS WAS THE ANALYTIC GEOMETRY. IT IS OFTEN REFERRED TO AS CARTESIAN GEOMETRY, WHICH IS A METHOD OF STUDYING GEOMETRY BY MEANS OF A COORDINATE SYSTEM AND ASSOCIATED ALGEBRA. IN ANALYTIC GEOMETRY, WE DESCRIBE PROPERTIES OF GEOMETRIC FIGURES SUCH AS CIRCLES, ETC., IN TERMS OF ORDERED PAIRS AND EQUATIONS.

4.1 DISTANCE BETWEEN TWO POINTS

IN **GRADE 9**, YOU HAVE DISCUSSED THE NUMBER LINE AND YOU HAVE SEEN THAT THERE IS A ONE-TO-ONE CORRESPONDENCE BETWEEN THE SET OF REAL NUMBERS AND THE SET OF POINTS ON A NUMBER LINE. YOU HAVE ALSO SEEN HOW TO LOCATE A POINT IN THE COORDINATE PLANE. DO YOU REMEMBER THE FACT THAT THERE IS A ONE-TO-ONE CORRESPONDENCE BETWEEN THE SET OF POINTS IN THE PLANE AND THE SET OF ALL ORDERED PAIRS OF REAL NUMBERS?

THE FOLLOWING ACTIVITY WILL HELP YOU TO REVIEW THE FACTS YOU DISCUSSED IN **GRADE 9**.

ACTIVITY 4.1



- 1 CONSIDER THE NUMBER LINE GIVEN IN **FIGURE 4.1**




Figure 4.1

- A** FIND THE COORDINATES OF POINTS **P, Q AND B**.
- B** FIND THE DISTANCE BETWEEN POINTS
- I** P AND Q **II** Q AND B **III** P AND B
- 2 ON A NUMBER LINE, THE TWO POINTS HAVE COORDINATES x_1 AND x_2 .
- A** FIND THE DISTANCE BETWEEN P AND Q .
- B** FIND THE DISTANCE BETWEEN Q AND B .
- C** DISCUSS THE RELATIONSHIP BETWEEN YOUR ANSWERS IN **A** AND **B**.
- D** DISCUSS THE RELATIONSHIP BETWEEN $|x_1 - x_2|$ AND $|x_2 - x_1|$.
- 3 HOW DO YOU PLOT THE COORDINATES OF POINTS IN THE COORDINATE PLANE?
- 4 WHAT ARE THE COORDINATES OF THE ORIGIN OF THE xy PLANE?
- 5 DRAW A COORDINATE PLANE AND PLOT THE FOLLOWING POINTS. $P(3, -4)$, $Q(-3, -2)$, $R(-2, 0)$, $S(4, 0)$, $T(2, 3)$, $U(-4, 5)$ AND $V(0, 0)$.

- 6 THE POSITION OF EACH POINT ON THE COORDINATE PLANE IS DETERMINED BY ITS PAIR OF NUMBERS.
- A WHAT IS THE COORDINATE OF A POINT ON THE y
 - B WHAT IS THE COORDINATE OF A POINT ON THE x
- 7 LET $P(2, 3)$ AND $Q(2, 8)$ BE POINTS ON THE COORDINATE PLANE.
- A PLOT THE POINTS.
 - B IS THE LINE THROUGH POINTS P AND Q VERTICAL OR HORIZONTAL?
 - C WHAT IS THE DISTANCE BETWEEN P AND Q ?
- 8 LET $R(-2, 4)$ AND $T(5, 4)$ BE POINTS ON THE COORDINATE PLANE.
- A PLOT THE POINTS.
 - B IS THE LINE THROUGH POINTS R AND T VERTICAL OR HORIZONTAL?
 - C WHAT IS THE DISTANCE BETWEEN POINTS R AND T ?

Distance between points in a plane

SUPPOSE $P(x_1, y_1)$ AND $Q(x_2, y_2)$ ARE TWO DISTINCT POINTS ON THE COORDINATE PLANE. WE CAN FIND THE DISTANCE BETWEEN THEM BY CONSIDERING THREE CASES.

Case i WHEN P AND Q ARE ON A LINE PARALLEL TO THE x -AXIS (THAT IS, \overline{PQ} IS A HORIZONTAL SEGMENT) AS IN FIGURE 4.2




Figure 4.2

SINCE THE TWO POINTS HAVE THE SAME y -COORDINATE, THE DISTANCE BETWEEN THEM IS

$$PQ = |x_2 - x_1|$$

Case ii WHEN P AND Q ARE ON A LINE PARALLEL TO THE y -AXIS (THAT IS, \overline{PQ} IS A VERTICAL SEGMENT) AS IN FIGURE 4.3




Figure 4.3

SINCE THE TWO POINTS HAVE THE SAME x -COORDINATE, THE DISTANCE BETWEEN THEM IS

$$PQ = |y_2 - y_1|$$

Case iii WHEN \overline{PQ} IS NEITHER VERTICAL NOR HORIZONTAL (THE GENERAL CASE).

TO FIND THE DISTANCE BETWEEN THE POINTS P AND Q , DRAW A LINE PASSING THROUGH P PARALLEL TO THE x -AXIS AND DRAW A LINE PASSING THROUGH Q PARALLEL TO THE y -AXIS. THE HORIZONTAL LINE AND THE VERTICAL LINE INTERSECT AT R .




Figure 4.4

USING CASE I AND CASE II, WE HAVE

$$PR = |x_2 - x_1| \text{ AND } RQ = |y_2 - y_1|$$

SINCE PRQ IS A RIGHT ANGLED TRIANGLE, WE CAN USE **Pythagoras' Theorem** TO FIND THE DISTANCE BETWEEN POINTS P AND Q AS FOLLOWS:

$$PQ^2 = PR^2 + RQ^2 = |x_2 - x_1|^2 + |y_2 - y_1|^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$$

$$\text{THEREFORE } PQ = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

THE RADICAL HAS POSITIVE SIGN (WHY?).

IN GENERAL, THE DISTANCE BETWEEN ANY TWO POINTS $P(x_1, y_1)$ AND $Q(x_2, y_2)$ IS GIVEN BY

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

THIS IS CALLED **distance formula**.

EXAMPLE 1 FIND THE DISTANCE BETWEEN THE GIVEN POINTS.

- A** $A(1, \sqrt{2})$ AND $B(1, \sqrt{2})$ **B** $P\left(\frac{17}{4}, -2\right)$ AND $Q\left(\frac{1}{4}, -2\right)$
C $R(-\sqrt{2}, -1)$ AND $S(\sqrt{2}, -\sqrt{2})$ **D** $A(a, -b)$ AND $B(-b, a)$

SOLUTION:

A $AB = d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$
 $= \sqrt{(1-1)^2 + (-\sqrt{2}-\sqrt{2})^2}$
 $= \sqrt{(0)^2 + (-2\sqrt{2})^2} = 2\sqrt{2}$.

OR, MORE SIMPLY

$$AB = |y_2 - y_1| = |-\sqrt{2} - \sqrt{2}| = 2\sqrt{2} \text{ UNIT}$$

$$\begin{aligned}
 \text{B } PQ = d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \\
 &= \sqrt{\left(\frac{1}{4} - \frac{17}{4}\right)^2 + (-2 - (-2))^2} \\
 &= \sqrt{\left(\frac{-16}{4}\right)^2 + (0)^2} = \sqrt{(-4)^2} = \sqrt{16} = 4 \text{ UNIT}
 \end{aligned}$$

OR, MORE SIMPLY

$$\begin{aligned}
 PQ &= |x_2 - x_1| = \left| \frac{1}{4} - \frac{17}{4} \right| \\
 &= 4 \text{ UNITS}
 \end{aligned}$$

$$\begin{aligned}
 \text{C } RS = d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(\sqrt{2} - (-\sqrt{2}))^2 + (-\sqrt{2} - (-1))^2} \\
 &= \sqrt{(2\sqrt{2})^2 + (1 - \sqrt{2})^2} = \sqrt{11 - 2\sqrt{2}}
 \end{aligned}$$

$$\begin{aligned}
 \text{D } AB = d &= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(-b - a)^2 + (a - (-b))^2} \\
 &= \sqrt{(b + a)^2 + (a + b)^2} = \sqrt{2(a + b)^2} = \sqrt{2}|a + b| \text{ UNIT}
 \end{aligned}$$

Exercise 4.1


- IN EACH OF THE FOLLOWING, FIND THE DISTANCE BETWEEN THE TWO GIVEN POINTS.

<p>A A (1, -5) AND B (7, 3)</p> <p>C E($\sqrt{2}$, 1) AND F($\sqrt{6}$, $\sqrt{3}$)</p> <p>E THE ORIGIN AND K($\frac{\sqrt{2}}{2}$, $\frac{-\sqrt{2}}{2}$)</p> <p>G P($\sqrt{2}$, $\sqrt{3}$) AND Q($\sqrt{2}$, $\sqrt{3}$)</p>	<p>B C($-2, \frac{1}{2}$) AND D($\frac{1}{2}, 2$)</p> <p>D G(a, -b) AND H(-a, b)</p> <p>F L($\sqrt{2}$, 1) AND M($\sqrt{2}$, 1)</p> <p>H R($\sqrt{2}a$, c) AND T($\sqrt{2}b$, c)</p>
--	--
- USING THE DISTANCE FORMULA, SHOW THAT THE DISTANCE BETWEEN P AND Q IS $|x_2 - x_1|$, WHEN \overline{PQ} IS HORIZONTAL AND $|y_2 - y_1|$, WHEN \overline{PQ} IS VERTICAL.
- LET A (3, -7) AND B (-1, 4) BE TWO ADJACENT VERTICES OF A SQUARE. CALCULATE THE SIDE LENGTH OF THE SQUARE.
- P (3, 5) AND Q (1, -3) ARE TWO OPPOSITE VERTICES OF A SQUARE. FIND ITS AREA.
- SHOW THAT THE PLANE FIGURE WITH VERTICES:

A A (5, -1), B (2, 3) AND C (1, 1) IS A RIGHT ANGLED TRIANGLE.	B A (-4, 3), B (4, -3) AND C ($3\sqrt{3}$, $4\sqrt{3}$) IS AN EQUILATERAL TRIANGLE.	C A (2, 3), B (6, 8), C (7, -1) IS AN ISOSCELES TRIANGLE.
---	---	--
- AN EQUILATERAL TRIANGLE HAS TWO VERTICES AT A (-4, 0) AND B (4, 0). WHAT COULD BE THE COORDINATES OF THE THIRD VERTEX?
- WHAT ARE THE POSSIBLE VALUES OF X IF POINT A(4, 4) IS 10 UNITS AWAY FROM B (0, -2)?

4.2 DIVISION OF A LINE SEGMENT

RECALL THAT, A LINE SEGMENT PASSING THROUGH TWO POINTS A AND B IS **horizontal** IF THE TWO POINTS HAVE THE SAME **ordinate**. I.E., A LINE SEGMENT WHOSE END-POINTS, A AND



$B(x_2, y)$ IS A HORIZONTAL LINE SEGMENT AS SHOWN IN **FIGURE 4.5**

What is the mid-point of \overline{AB} ?

Figure 4.5

ACTIVITY 4.2

- 1 DEFINE THE RATIO OF TWO QUANTITIES.
- 2 WHAT IS MEANT BY THE RATIO OF THE LENGTH OF TWO LINE SEGMENTS?
- 3 IN **FIGURE 4.6** FIND THE RATIO OF THE LENGTHS OF \overline{AP} AND \overline{PB} .




Figure 4.6

- 4 WHAT IS MEANT BY A POINT DIVIDES A LINE SEGMENT INTERNALLY?
- 5 PLOT THE FOLLOWING POINTS ON THE COORDINATE PLANE AND FIND THE MID-POINT OF THE LINE SEGMENT JOINING THE POINTS.

A $A(2, -1)$ AND $B(2, 5)$ **B** $C(-3, 3)$ AND $D(3, 3)$ **C** $E(2, 0)$ AND $F(-2, 4)$.

CONSIDER THE HORIZONTAL LINE SEGMENT WITH END-POINTS A AND B AS SHOWN IN **FIGURE 4.7** IN TERMS OF THE COORDINATES, DETERMINE THE COORDINATES OF THE POINT $P(x_0, y_0)$ THAT DIVIDES \overline{AB} INTERNALLY IN THE RATIO

CLEARLY, THE RATIO OF THE LINE SEGMENTS IS GIVEN BY $\frac{AP}{PB}$

THE DISTANCE BETWEEN A AND P IS $AP = x_0 - x_1$.

THE DISTANCE BETWEEN P AND B IS $PB = x_2 - x_0$.

THEREFORE $\frac{AP}{PB} = \frac{m}{n}$ I.E., $\frac{x_0 - x_1}{x_2 - x_0} = \frac{m}{n}$.

SOLVING THIS EQUATION FOR

$$\Rightarrow n(x_0 - x_1) = m(x_2 - x_0)$$

$$\Rightarrow nx_0 - nx_1 = mx_2 - mx_0$$

$$\Rightarrow nx_0 + mx_0 = nx_1 + mx_2$$




Figure 4.7

$$\Rightarrow x_0(n + m) = nx_1 + mx_2$$

$$\Rightarrow x_0 = \frac{nx_1 + mx_2}{n + m}$$

SINCE \overline{AB} IS PARALLEL TO THE X-AXIS (AS \overline{AB} IS A HORIZONTAL LINE SEGMENT) AND OBVIOUSLY, $y_0 = y$, THEREFORE, THE POINT IS $\left(\frac{nx_1 + mx_2}{n + m}, y\right)$.

GIVEN A LINE SEGMENT WITH END POINT COORDINATES $P(x_1, y_1)$ AND $Q(x_2, y_2)$, LET US FIND THE COORDINATES OF THE POINT DIVIDING THE LINE SEGMENT INTERNALLY IN THE RATIO I.E., $\frac{PR}{RQ} = \frac{m}{n}$, WHERE m AND n ARE GIVEN POSITIVE REAL NUMBERS.




Figure 4.8

LET THE COORDINATES OF R BE (x_0, y_0) . ASSUME THAT x_2 AND $x_1 \neq y_2$.

IF YOU DRAW LINES THROUGH THE POINTS PARALLEL TO THE AXES AS SHOWN IN FIGURE 4.8 THE POINTS S AND T HAVE THE COORDINATES (x_0, y_1) AND (x_2, y_0) , RESPECTIVELY.

$$PS = x_0 - x_1, RT = x_2 - x_0, SR = y_0 - y_1 \text{ AND } QT = y_2 - y_0$$

SINCE TRIANGLES $\triangle PSR$ AND $\triangle RTQ$ ARE SIMILAR (WHY?),

$$\frac{PS}{RT} = \frac{PR}{RQ} \text{ AND } \frac{SR}{TQ} = \frac{PR}{RQ}$$

$$\frac{x_0 - x_1}{x_2 - x_0} = \frac{m}{n} \text{ AND } \frac{y_0 - y_1}{y_2 - y_0} = \frac{m}{n}$$

SOLVING FOR x_0 AND y_0

$$\Rightarrow n(x_0 - x_1) = m(x_2 - x_0) \text{ AND } n(y_0 - y_1) = m(y_2 - y_0)$$

$$\Rightarrow nx_0 - nx_1 = mx_2 - mx_0 \text{ AND } ny_0 - ny_1 = my_2 - my_0$$

$$\Rightarrow nx_0 + mx_0 = nx_1 + mx_2 \text{ AND } ny_0 + my_0 = ny_1 + my_2$$

$$\Rightarrow x_0(n + m) = nx_1 + mx_2 \text{ AND } y_0(n + m) = ny_1 + my_2$$

$$\Rightarrow x_0 = \frac{nx_1 + mx_2}{n + m} \text{ AND } y_0 = \frac{ny_1 + my_2}{n + m}$$

THE POINT $R(x_0, y_0)$ DIVIDING THE LINE SEGMENT INTERNALLY IN THE RATIO $m:n$ IS GIVEN BY

$$R(x_0, y_0) = \left(\frac{nx_1 + mx_2}{n+m}, \frac{ny_1 + my_2}{n+m} \right)$$

THIS IS CALLED SECTION FORMULA.

EXAMPLE 1 FIND THE COORDINATES OF THE POINTS THAT DIVIDE THE LINE SEGMENT WITH END-POINTS A (6, 2) AND B (1, -4) IN THE RATIO 2:3.

SOLUTION: PUT $(x_1, y_1) = (6, 2)$, $(x_2, y_2) = (1, -4)$, $m = 2$ AND $n = 3$. USING THE SECTION FORMULA, YOU HAVE

$$\begin{aligned} R(x_0, y_0) &= \left(\frac{nx_1 + mx_2}{n+m}, \frac{ny_1 + my_2}{n+m} \right) = \left(\frac{3 \times 6 + 2 \times 1}{3+2}, \frac{3 \times 2 + 2 \times (-4)}{3+2} \right) \\ &= \left(\frac{18+2}{5}, \frac{6-8}{5} \right) = \left(4, -\frac{2}{5} \right) \\ &\text{THEREFORE, } R \left(4, -\frac{2}{5} \right). \end{aligned}$$

EXAMPLE 2 A LINE SEGMENT HAS END-POINTS (-2, -3) AND (7, 12) AND IT IS DIVIDED INTO THREE EQUAL PARTS. FIND THE COORDINATES OF THE POINTS THAT TRISECT THE SEGMENT.

SOLUTION: THE FIRST POINT DIVIDES THE LINE SEGMENT IN THE RATIO 1:2, AND HENCE

$$\begin{aligned} x_0 &= \frac{nx_1 + mx_2}{n+m} \text{ AND } y_0 = \frac{ny_1 + my_2}{n+m} \\ \text{SO, } x_0 &= \frac{2 \times (-2) + 1 \times 7}{1+2} \text{ AND } y_0 = \frac{2 \times (-3) + 1 \times 12}{1+2} \\ \Rightarrow x_0 &= \frac{-4+7}{3} \text{ AND } y_0 = \frac{-6+12}{3} \Rightarrow x_0 = 1 \text{ AND } y_0 = 2 \end{aligned}$$

THEREFORE, THE FIRST POINT IS (1, 2).

THE SECOND POINT DIVIDES THE LINE SEGMENT IN THE RATIO 2:1. THUS,

$$\begin{aligned} x_0 &= \frac{nx_1 + mx_2}{n+m} \text{ AND } y_0 = \frac{ny_1 + my_2}{n+m} \\ \text{SO, } x_0 &= \frac{1 \times (-2) + 2 \times 7}{1+2} \text{ AND } y_0 = \frac{1 \times (-3) + 2 \times 12}{1+2} \\ \Rightarrow x_0 &= \frac{-2+14}{3} \text{ AND } y_0 = \frac{-3+24}{3} \\ \Rightarrow x_0 &= 4 \text{ AND } y_0 = 7. \end{aligned}$$

THEREFORE, THE SECOND POINT IS (4, 7).

The mid-point formula

A POINT THAT DIVIDES A LINE SEGMENT INTO TWO EQUAL PARTS IS THE MID-POINT OF THE

ACTIVITY 4.3



- 1 CONSIDER THE POINTS P (2, 1) AND Q (12, 1).
 - A FIND THE DISTANCE BETWEEN P AND Q
 - B IF R IS A POINT WITH COORDINATES (7, 1),
 - I FIND PR II FIND RQ
 - III IS PR EQUAL TO RQ IV WHAT IS THE MID-POINT OF PQ
 - C DIVIDE \overline{PQ} IN THE RATIO 1:1.
- 2 FIND THE COORDINATES OF THE MID-POINT OF EACH OF THE FOLLOWING LINE SEGMENTS WITH GIVEN END-POINTS:
 - I P (x_1, y_1) AND Q (x_2, y_2). II R (x_1, y_1) AND S (x_2, y_1).

WHICH OF THE ABOVE SEGMENTS ARE HORIZONTAL?

LET P (x_1, y_1) AND Q (x_2, y_2) BE THE END-POINTS OF

IF $PR = RQ$ (THE CASE WHERE R IS THE MID-POINT OF THE LINE SEGMENT PQ)

NOW LET US DERIVE THE MID-POINT FORMULA.

$$\begin{aligned}
 R(x, y) &= \left(\frac{nx_1 + mx_2}{n+m}, \frac{ny_1 + my_2}{n+m} \right) \\
 &= \left(\frac{nx_1 + nx_2}{n+n}, \frac{ny_1 + ny_2}{n+n} \right) = \left(\frac{n(x_1 + x_2)}{2n}, \frac{n(y_1 + y_2)}{2n} \right) \quad (\text{As } m = n) \\
 &= \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)
 \end{aligned}$$

THIS IS THE FORMULA USED TO FIND THE MID-POINT OF THE LINE SEGMENT WHOSE END POINTS ARE P (x_1, y_1) AND Q (x_2, y_2).

THE MID-POINT OF THE LINE SEGMENT JOINING THE POINTS (x_1, y_1) AND (x_2, y_2) IS GIVEN BY

$$M(x, y) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

EXAMPLE 3 FIND THE COORDINATES OF THE MID-POINT OF THE FOLLOWING LINE SEGMENTS

- A P (-3, 2) AND Q (5, -4)
- B P ($3 - \sqrt{2}, 3 + \sqrt{2}$) AND Q ($1 + \sqrt{2}, 3 - \sqrt{2}$).

SOLUTION:

A $M(x_0, y_0) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

$x_0 = \frac{x_1 + x_2}{2}$ AND $y_0 = \frac{y_1 + y_2}{2}$

$x_0 = \frac{-3 + 5}{2} = 1$ AND $y_0 = \frac{2 - 4}{2} = -1$

THEREFORE $M(x_0, y_0) = (1, -1)$.

B $M(x_0, y_0) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

$x_0 = \frac{x_1 + x_2}{2}$ AND $y_0 = \frac{y_1 + y_2}{2}$

$x_0 = \frac{3 - \sqrt{2} + 1 + \sqrt{2}}{2}$ AND $y_0 = \frac{3 + \sqrt{2} + 3 - \sqrt{2}}{2}$

$x_0 = \frac{4}{2} = 2$ AND $y_0 = \frac{6}{2} = 3$

THEREFORE $M(x_0, y_0) = (2, 3)$.

Group Work 4.1



- 1** A LINE SEGMENT HAS END-POINTS P (-3, 1) AND Q (4, 3).
 - A** WHAT IS THE LENGTH OF THE LINE SEGMENT?
 - B** FIND THE COORDINATES OF THE MID-POINT OF THE SEGMENT.
- 2** A LINE SEGMENT HAS ONE END-POINT AT A (4, 3). IF ITS MID-POINT IS AT M (1, -1), WHERE IS THE OTHER END-POINT?
- 3** FIND THE POINTS THAT DIVIDE THE LINE SEGMENT WITH END-POINTS AT P Q (-6, 7) INTO THREE EQUAL PARTS.
- 4** LET A (-2, -1), B (6, -1), C (6, 3) AND D (-2, 3) BE VERTICES OF A RECTANGLE. SUPPOSE P, Q, R AND S ARE MID-POINTS OF THE SIDES OF THE RECTANGLE.
 - I** WHAT IS THE AREA OF RECTANGLE ABCD?
 - II** WHAT IS THE AREA OF QUADRILATERAL PQRS?
 - III** GIVE THE RATIO OF THE AREAS IN I AND II.

Exercise 4.2

- 1 FIND THE COORDINATES OF THE MID-POINT OF THE LINE SEGMENTS JOINING THE POINTS

A A (1, 4) AND B(-2, 2)	B (a, b) AND THE ORIGIN
C M (p, q) AND N(q, p)	D $(\frac{1}{2}, -1)$ AND $(-\frac{5}{2}, 1)$
E E(1+√2, √2) AND F(2√2, 2√2)	F G(√5, 1-√3) AND H(√5, 1+√3)
- 2 THE MID-POINT OF A LINE SEGMENT IS (2, 5). ONE END-POINT OF THE SEGMENT IS P (1, -3). FIND THE COORDINATES OF THE OTHER END-POINT.
- 3 FIND THE COORDINATES OF THE POINT WHICH BISECTS THE LINE SEGMENT JOINING THE POINTS A (1, 3) AND B(-4, -3) IN THE RATIO 2:3.
- 4 A LINE SEGMENT HAS END-POINTS P(1, 2) AND Q (5, 2). FIND THE COORDINATES OF THE POINTS THAT TRISECT THE SEGMENT.
- 5 FIND THE MID-POINTS OF THE SIDES OF THE TRIANGLE WITH VER- TICES A (1, 2), B (4, 6) AND C (3, -1).

4.3 EQUATION OF A LINE

4.3.1 Gradient (slope) of a Line

FROM YOUR EVERYDAY EXPERIENCE, YOU MIGHT BE FAMILIAR WITH THE IDEA OF GRADIENT (SLOPE). A hill may be steep or may rise very slowly. THE NUMBER THAT DESCRIBES THE STEEPNESS OF A HILL IS CALLED THE GRADIENT (SLOPE) OF THE HILL.



WE MEASURE THE GRADIENT OF A HILL BY THE RATIO OF THE VERTICAL RISE TO THE HORIZONTAL RUN.

Figure 4.9


ACTIVITY 4.4

GIVEN POINTS P (1, 2), Q (-4), R (0, -1) AND S (3, 8)

- A** FIND THE VALUE OF $\frac{y_2 - y_1}{x_2 - x_1}$ TAKING
- I** P AND Q **II** P AND R **III** Q AND R **IV** R AND S
- B** ARE THE VALUES OBTAINED ABOVE EQUAL? WHAT DO YOU CALL THESE VALUES?



IN COORDINATE GEOMETRY, THE GRADIENT OF A NON-VERTICAL STRAIGHT LINE IS THE RATIO OF "CHANGE IN y-COORDINATES" TO THE CORRESPONDING "CHANGE IN x-COORDINATES". THAT IS, THE SLOPE OF A LINE THROUGH P AND Q IS THE RATIO OF THE VERTICAL DISTANCE FROM P TO Q TO THE HORIZONTAL DISTANCE FROM P TO Q .



IF WE DENOTE THE GRADIENT OF A LINE BY THE LETTER m , THEN

$$m = \frac{\text{change in } y\text{-coordinates}}{\text{change in } x\text{-coordinates}} = \frac{y_2 - y_1}{x_2 - x_1}; x_1 \neq x_2$$


Figure 4.10

Definition 4.1

If (x_1, y_1) and (x_2, y_2) are points on a line with $x_1 \neq x_2$, then the **gradient** of the line, denoted by m , is given by

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

ACTIVITY 4.5



- 1 IF A (x_1, y_1) AND $B(x_2, y_2)$ ARE DISTINCT POINTS ON A LINE WITH $x_1 = x_2$, THEN WHAT CAN BE SAID ABOUT THE GRADIENT OF THE LINE? IS THE LINE VERTICAL OR HORIZONTAL?
- 2 WHAT IS THE GRADIENT OF ANY HORIZONTAL LINE?
- 3 CONSIDER THE LINE WITH EQUATION $y = 2x + 3$. TAKE THREE DISTINCT POINTS P_1, P_2 AND P_3 ON THE LINE.
 - A FIND THE GRADIENT OF THE LINE TAKING P_1 AND P_2 AS POINTS.
 - B FIND THE GRADIENT OF THE LINE TAKING P_1 AND P_3 AS POINTS.
 - C WHAT DO YOU OBSERVE FROM A AND B?
- 4 LET P_1, P_2, P_3 AND P_4 BE POINTS ON A NON-VERTICAL STRAIGHT LINE WITH COORDINATES $(x_1, y_1), (x_2, y_2), (x_3, y_3)$ AND (x_4, y_4) RESPECTIVELY. FIND:
 - A THE GRADIENT OF THE LINE TAKING P_1 AND P_2 AS POINTS.
 - B THE GRADIENT OF THE LINE TAKING P_1 AND P_3 AS POINTS.
 - C ARE THE RATIOS $\frac{y_2 - y_1}{x_2 - x_1}$ AND $\frac{y_4 - y_3}{x_4 - x_3}$ EQUAL?
 - D COULD YOU CONCLUDE THAT THE GRADIENT OF A LINE DOES NOT DEPEND ON THE CHOICE OF POINTS ON THE LINE?

EXAMPLE 1 FIND THE GRADIENT OF THE LINE PASSING THROUGH EACH OF THE FOLLOWING OF POINTS:

A P (-7, 2) AND Q (4, 3)

B A ($\sqrt{2}$, 1) AND B ($-\sqrt{2}$, -3)

C P (2, -3) AND Q (5, -3)

D A ($-\frac{1}{2}$, -2) AND B ($-\frac{1}{2}$, 2)

SOLUTION:

A $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{3 - 2}{4 - (-7)} = \frac{1}{11}$

B $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - 1}{-\sqrt{2} - \sqrt{2}} = \frac{-4}{-2\sqrt{2}} = \frac{2}{\sqrt{2}} = \sqrt{2}$

C $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-3)}{5 - 2} = \frac{-3 + 3}{3} = \frac{0}{3} = 0$

SO, $m = 0$. IS THE LINE HORIZONTAL? WHAT IS ITS EQUATION?

D $x_1 = -\frac{1}{2}$ AND $x_2 = -\frac{1}{2}$

THE LINE IS VERTICAL. SO IT HAS NO MEASURABLE GRADIENT.

THE EQUATION OF THE LINE IS $x = -\frac{1}{2}$ OR SIMPLY $x = -\frac{1}{2}$

Note: GRADIENT FOR A VERTICAL LINE IS NOT DEFINED.

EXAMPLE 2 CHECK THAT THE LINES THROUGH P (0, 1) AND Q (-1, 4) AND THROUGH

R ($\frac{2}{3}$, 0) AND T (1, -1) HAVE SAME GRADIENTS. ARE THE LINES PARALLEL?

SOLUTION: FOR ℓ , $m_1 = \frac{4 - 1}{-1 - 0} = \frac{3}{-1} = -3$. FOR ℓ , $m_2 = \frac{-1 - 0}{1 - \frac{2}{3}} = \frac{-1}{\frac{1}{3}} = -3$.

HERE $m_1 = m_2$. DRAW THE LINES AND SEE IF PARALLEL TO ℓ

Exercise 4.3

1 FIND THE GRADIENTS OF THE LINES PASSING THROUGH THE FOLLOWING POINTS:

A A (4, 3) AND B (8, 11)

B P (3, 7) AND Q (1, 9)

C C ($\sqrt{2}$, -9) AND D ($2\sqrt{2}$, -7)

D R (-5, -2) AND S (7, -8)

E E (5, 8) AND F (-2, 8)

F H (1, 7) AND K (-1, -6)

G R (1, b) AND S (a, b), $b \neq 1$.

- 2 A (2, -3), B (7, 5) AND C(-2, 9) ARE THE VERTICES OF TRIANGLE ABC. FIND THE GRADIENT OF EACH OF THE SIDES OF THE TRIANGLE.
- 3 GIVEN THREE POINTS, P(5), Q (1, -2) AND R (5, 4), FIND THE GRADIENT OF \overline{PQ} AND \overline{QR} . WHAT DO YOU CONCLUDE FROM YOUR RESULT?
- 4 USE GRADIENTS TO SHOW THAT THE POINTS P(12) AND Q(-7, 0) ARE COLLINEAR, I.E., ALL LIE ON THE SAME STRAIGHT LINE.
- 5 SHOW THAT THE LINE PASSING THROUGH THE POINTS A(0, 3) AND B(2, 0), ALSO PASSES THROUGH THE POINT C(4, -3).

4.3.2 Slope of a Line in Terms of Angle of Inclination

THE ANGLE MEASURED FROM THE POSITIVE x-AXIS TO A LINE, IN ANTICLOCKWISE DIRECTION, IS CALLED THE **inclination of the line** OR THE ANGLE OF INCLINATION OF THE LINE. THIS ANGLE IS ALWAYS LESS THAN 180°




Figure 4.11

Group Work 4.2



CONSIDER THE RIGHT ANGLED TRIANGLE **Figure 4.12**

- 1 HOW LONG IS THE HYPOTENUSE
- 2 WHAT IS TANGENT OF ANGLE
- 3 WHAT IS MEASURE OF ANGLE BO
- 4 WHAT IS THE ANGLE OF INCLINATION OF LINE l
- 5 WHAT IS THE TANGENT OF THE ANGLE OF INCLINATION?
- 6 BY FINDING THE COORDINATES OF A, CALCULATE THE SLOPE OF LINE l




Figure 4.12

- 7 WHAT RELATIONSHIP DO YOU SEE BETWEEN YOUR ANSWERS AND ABOVE QUESTIONS 5

THE ABOVE GROUP WILL HELP YOU TO UNDERSTAND THE RELATIONSHIP BETWEEN SLOPE AND ANGLE OF INCLINATION.

FOR A NON-VERTICAL LINE, THIS ANGLE IS THE **slope** OF THE LINE. OBSERVE THE FOLLOWING.




Figure 4.13

IN FIGURE 4.13A ABOVE, AS y_1 REPRESENTS THE DISTANCE $-x_1$ REPRESENTS THE DISTANCE PR, THE SLOPE OF THE STRAIGHT LINE IS REPRESENTED BY THE RATIO

$$m = \frac{RQ}{PR} = \frac{y_2 - y_1}{x_2 - x_1} = \text{TAN}(\angle RPQ)$$

$$\therefore m = \text{TAN}$$

A LINE MAKING AN ACUTE ANGLE OF WITH THE POSITIVE DIRECTION OF THE x HAS POSITIVE SLOPE.

SIMILARLY, A LINE WITH OBTUSE ANGLE OF INCLINATION HAS NEGATIVE SLOPE.

$$\text{SLOPE OF } \ell = \frac{RQ}{PR} = \frac{y_2 - y_1}{x_1 - x_2} = -\frac{y_2 - y_1}{x_2 - x_1} = -\text{TAN}(180^\circ - \theta) \Rightarrow -(\text{TAN } \theta)$$

(In Unit 5, this will be clarified)

ACTIVITY 4.6

- 1 HOW WOULD YOU DESCRIBE THE LINE PASSING THROUGH THE POINTS WITH COORDINATES (x_1, y_1) AND (x_2, y_2) ? IS IT PERPENDICULAR TO THE x -AXIS OR THE y -AXIS? WHAT IS THE TANGENT OF THE ANGLE BETWEEN THIS LINE AND THE x -AXIS?
- 2 SUPPOSE A LINE PASSES THROUGH THE POINTS WITH COORDINATES (x_1, y_1) AND (x_2, y_2) . FIND THE TANGENT OF THE ANGLE FORMED BY THE LINE WITH THE x -AXIS. WHAT IS THE SLOPE OF THIS LINE?
- 3 WHAT IS THE ANGLE OF INCLINATION OF A LINE WITH THE x -AXIS?



IN GENERAL, THE SLOPE OF A LINE MAY BE EXPRESSED IN TERMS OF THE COORDINATES (x_1, y_1) AND (x_2, y_2) ON THE LINE AS FOLLOWS:

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \tan \theta, \quad x_2 \neq x_1$$

WHERE θ IS THE ANTICLOCKWISE ANGLE BETWEEN THE POSITIVE x -AXIS AND THE LINE.

EXAMPLE 3 FIND THE SLOPE OF A LINE, IF ITS INCLINATION IS:

- A** 60° **B** 135°

SOLUTION:

A SLOPE : $m = \tan 60^\circ = \sqrt{3}$

B SLOPE : $m = \tan 135^\circ = \tan (180^\circ - 45^\circ) = -\tan 45^\circ = -1$

Note: IF θ IS AN OBTUSE ANGLE, THEN $\tan \theta = \tan (180^\circ - \theta)$.

EXAMPLE 4 FIND THE ANGLE OF INCLINATION OF THE LINE

- A** CONTAINING THE POINTS A(3, -3) AND B(-1, 1)
B CONTAINING THE POINTS C(0, 5) AND D(4, 5).

SOLUTION:

A $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - (-3)}{-1 - 3} = -1$. SO $\tan \theta = -1$ AND HENCE $\theta = 135^\circ$.

B $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 5}{4 - 0} = 0$, $\tan \theta = 0$. SO, $\theta = 0^\circ$.

Note: LET m BE THE SLOPE OF A NON-VERTICAL LINE.

- I** IF $m > 0$, THEN THE LINE RISES FROM LEFT TO RIGHT AS SHOWN IN FIGURE 4.14A.
- II** IF $m < 0$, THEN THE LINE FALLS FROM LEFT TO RIGHT AS SHOWN IN FIGURE 4.14B.
- III** IF $m = 0$, THEN THE LINE IS HORIZONTAL AS IN FIGURE 4.14C.




Figure 4.14

Exercise 4.4

- 1 FIND THE SLOPE OF THE LINE WHOSE ANGLE OF INCLINATION IS:
A 30° **B** 75° **C** 150° **D** 90° **E** 0°
- 2 FIND THE ANGLE OF INCLINATION OF THE LINE IF ITS SLOPE IS:
A $-\sqrt{3}$ **B** $\frac{-\sqrt{3}}{3}$ **C** 1 **D** $\frac{1}{\sqrt{3}}$ **E** 0.
- 3 THE POINTS A(0, 0), B(0, 2) AND C(2, 0) ARE VERTICES OF A TRIANGLE. FIND THE MEASURE OF THE THREE ANGLES OF THE TRIANGLE.

4.3.3 Different Forms of Equations of a Line

FROM EUCLIDEAN GEOMETRY, YOU MAY RECALL THAT THERE IS A UNIQUE LINE PASSING THROUGH TWO DISTINCT POINTS. THE EQUATION OF A LINE IS AN EQUATION WHICH IS SATISFIED BY THE COORDINATES OF EVERY POINT ON THE LINE AND IS NOT SATISFIED BY THE COORDINATES OF ANY POINT NOT ON THE LINE.

THE EQUATION OF A STRAIGHT LINE CAN BE EXPRESSED IN DIFFERENT FORMS. SOME OF THEM ARE THE POINT-SLOPE FORM, THE SLOPE-INTERCEPT FORM AND THE TWO-POINT FORM.

ACTIVITY 4.7

- 1 SHOW THAT THE GRAPH OF THE EQUATION $x = 2$ CONTAINS POINTS A(2, 0), B(2, -1), C(2, 2) AND D(2, $\frac{1}{3}$).
- 2 CONSIDER THE GRAPH OF THE STRAIGHT LINE $x + 2y = 4$. DETERMINE WHICH OF THE FOLLOWING POINTS LIE ON THE LINE.
 A(3, -1), B(-1, 0), C($\frac{-1}{2}$, $\frac{3}{2}$), D(0, 1), E($\frac{-1}{2}$, 1), F(-2, -1) AND G(-4, 2)
- 3 WHICH OF THE FOLLOWING POINTS LIE ON THE LINE $y = 3x + 12$?
 A(-1, 9), B(-2, 12), C(0, 4), D($\frac{2}{5}$, 2), E(3, -10).
- 4 WHAT DO YOU CALL THE NUMBER b IN THE EQUATION $y = mx + b$ THAT POINT P(0, b)?
- 5 CONSIDER THE GRAPH OF THE STRAIGHT LINE $y = 2x + 3$. FIND ITS y -INTERCEPT AND x -INTERCEPT.
- 6 GIVE THE EQUATIONS OF THE LINES THROUGH THE POINTS:
A P(-1, 3) AND Q(4, 3) **B** R(-1, 1) AND S(4, -1).



The point-slope form of equation of a line

WE NORMALLY USE THIS FORM OF THE EQUATION OF A ~~DEFINITION~~ LINE ~~AND~~ SLOPE ~~IF~~ THE COORDINATES OF A POINT ON IT ARE GIVEN.




Figure 4.15

SUPPOSE YOU ARE ASKED TO FIND THE EQUATION OF THE STRAIGHT LINE WITH SLOPE 3 THROUGH THE POINT WITH COORDINATE (2, 3).

TAKE ~~IT~~ ~~TO~~ BE THE POINT (2, 3) AND, ~~LET~~ ~~BE~~ ANY OTHER POINT ON THE LINE AS SHOWN IN ~~FIGURE~~ 4.15 WHAT IS THE SLOPE OF THE STRAIGHT LINE JOINING THE POINTS WITH COORDINATES (x_1, y_1) AND (x_2, y_2) ?

WHAT IS THE SLOPE ~~IF~~ YOU ARE GIVEN THAT THE SLOPE OF THIS LINE IS 3. IF YOU HAVE ANSWERED CORRECTLY, YOU SHOULD OBTAIN

$$y = 3x - 3;$$

WHICH IS THE REQUIRED EQUATION OF THE STRAIGHT LINE.

IN GENERAL, SUPPOSE YOU WANT TO FIND THE EQUATION OF THE STRAIGHT LINE WITH SLOPE m THROUGH THE POINT WITH COORDINATES (x_1, y_1) WHICH HAS SLOPE m . TAKE THE POINT WITH GIVEN COORDINATES ~~AS~~ ~~BE~~ TAKE ANY OTHER POINT ON THE LINE, ~~LET~~ ~~SAY~~ COORDINATES (x, y) AS SHOWN IN ~~FIGURE~~ 4.16




Figure 4.16

THEN THE SLOPE IS $\frac{y - y_1}{x - x_1}$

$$\Rightarrow y - y_1 = m(x - x_1) \text{ WHICH IS THE SAME AS } y = m(x - x_1) + y_1$$

THIS EQUATION IS CALLED THE **point-slope form** of the equation of a line

EXAMPLE 5 FIND THE EQUATION OF THE STRAIGHT LINE WHICH PASSES THROUGH THE POINT $(-3, 2)$

SOLUTION: ASSUME THAT THE POINT ANY POINT ON THE LINE OTHER THAN $(-3, 2)$.
 THUS, USING THE EQUATION $y - y_1 = m(x - x_1)$

$$\Rightarrow y - 2 = \frac{-3}{2}(x + 3)$$

$$\Rightarrow y = -\frac{3}{2}x - \frac{5}{2} \text{ OR } 2y + 3x + 5 = 0.$$

The slope-intercept form of equation of a line

CONSIDER THE EQUATION $y = mx + b$. WHEN $x = 0$, $y = b$. ALSO, WHEN $x = 1$, $y = m + b$ AS SHOWN IN

FIGURE 4.17

YOU CAN SEE THAT P $(0, b)$ IS THE POINT WHERE THE LINE WITH EQUATION $y = mx + b$ CROSSES THE y-AXIS. b IS CALLED THE **y-intercept** OF THE LINE).

LET Q BE $(1, m + b)$.

USING THE COORDINATES OF P AND Q, SHOW THAT THE SLOPE OF THE STRAIGHT LINE PASSING THROUGH P IS m




Figure 4.17

WRITING THE EQUATION OF THIS LINE THROUGH THE POINT $(0, b)$ WITH SLOPE m , USING THE POINT-SLOPE FORM, GIVES

$$y - b = m(x - 0) \Rightarrow y = mx + b$$

WHERE m IS SLOPE OF THE LINE AND b IS THE y-INTERCEPT OF THE LINE.

THIS EQUATION IS CALLED THE **slope-intercept form** OF THE EQUATION OF A LINE.

Note: THE SLOPE-INTERCEPT FORM OF EQUATION OF A LINE ENABLES US TO FIND THE y-INTERCEPT, ONCE THE EQUATION IS GIVEN.

EXAMPLE 6 FIND THE EQUATION OF THE LINE WITH SLOPE $\frac{2}{3}$ AND Y-INTERCEPT 3.

SOLUTION: HERE, $m = \frac{2}{3}$ AND THE Y-INTERCEPT IS 3.

THEREFORE, THE EQUATION OF THE LINE IS $y = \frac{2}{3}x + 3$

The two-point form of equation of a line

FINALLY, LET US LOOK AT THE SITUATION WHERE THE SLOPE IS NOT GIVEN BUT TWO POINTS ON THE LINE ARE GIVEN.

CONSIDER A STRAIGHT LINE WHICH PASSES THROUGH THE POINTS $P(x_1, y_1)$ AND $Q(x_2, y_2)$. IF $R(x, y)$ IS ANY POINT ON THE LINE OTHER THAN P OR Q , THEN THE SLOPE OF \overline{PR} IS

$$m = \frac{y - y_1}{x - x_1}, x \neq x_1$$

AND THE SLOPE OF \overline{PQ} IS

$$m = \frac{y_2 - y_1}{x_2 - x_1}, x_1 \neq x_2$$

BUT THE SLOPE OF \overline{PR} IS EQUAL TO THE SLOPE OF \overline{PQ}

$$\therefore \frac{y - y_1}{x - x_1} = \frac{y_2 - y_1}{x_2 - x_1}$$




Figure 4.18

THIS EQUATION IS CALLED THE **two-point form** of the equation of a line.

EXAMPLE 7 FIND THE EQUATION OF THE LINE PASSING THROUGH THE POINTS $P(-1, 5)$ AND $Q(3, 13)$.

SOLUTION: TAKING $(-1, 5)$ AS (x_1, y_1) AND $(3, 13)$ AS (x_2, y_2) , USE THE TWO-POINT FORM TO GET THE EQUATION OF THE LINE TO BE

$$y - 5 = \frac{13 - 5}{3 - (-1)}(x + 1) = 2x + 7 \text{ WHICH IMPLIES } 2x - y + 12 = 0$$

The general equation of a line

A FIRST DEGREE (LINEAR) EQUATION IN TWO VARIABLES IS AN EQUATION OF THE FORM;

$$Ax + By + C = 0$$

WHERE A AND B ARE FIXED REAL NUMBERS SUCH THAT $A^2 + B^2 \neq 0$

ALL THE DIFFERENT FORMS OF EQUATIONS OF LINES DISCUSSED ABOVE CAN BE EXPRESSED IN THE GENERAL FORM

$$Ax + By + C = 0$$

CONVERSELY, ONE CAN SHOW THAT ANY LINE IS THE EQUATION OF A LINE. SUPPOSE A LINE IS GIVEN AS

$$Ax + By + C = 0.$$

IF $B \neq 0$, THEN THE EQUATION MAY BE SOLVED AS FOLLOWS:

$$Ax + By + C = 0$$

$$By = -Ax - C$$

$$y = \frac{-A}{B}x - \frac{C}{B}$$

THIS EQUATION IS OF THE FORM $y = mx + c$ AND THEREFORE REPRESENTS A STRAIGHT LINE WITH SLOPE $m = -\frac{A}{B}$ AND INTERCEPT $c = -\frac{C}{B}$.

WHAT WILL BE THE EQUATION $Ax + By + C = 0$, IF $B \neq 0$ AND $A \neq 0$?

EXAMPLE 8 FIND THE SLOPE AND INTERCEPT OF THE LINE WHOSE GENERAL EQUATION IS $3x - 6y - 4 = 0$.

SOLUTION: SOLVING FOR y IN THE EQUATION $3x - 6y - 4 = 0$ GIVES,

$$-6y = -3x + 4 \Rightarrow y = \frac{-3x}{-6} + \frac{4}{-6} = \frac{1}{2}x - \frac{2}{3}$$

SO, THE SLOPE IS $\frac{1}{2}$ AND THE INTERCEPT IS $-\frac{2}{3}$.

EXAMPLE 9 WHAT IS THE EQUATION OF THE LINE PASSING THROUGH $(-2, 0)$ AND $(0, 5)$?

SOLUTION: USING TWO-POINT FORM:

$$y - 0 = \frac{5 - 0}{0 - (-2)}(x + 2)$$

WHICH GIVES US $2y + 10 = 0$ AS THE EQUATION OF THE LINE.

Exercise 4.5

1 FIND THE EQUATION OF THE LINE PASSING THROUGH THE GIVEN POINTS.

- | | | | |
|----------|------------------------------|----------|---|
| A | A $(-2, -4)$ AND B $(-1, 5)$ | B | C $(2, -4)$ AND D $(-1, 5)$ |
| C | E $(3, 7)$ AND F $(8, 7)$ | D | G $(1, 1)$ AND H $(1 + \sqrt{2}, 1 - \sqrt{2})$ |
| E | P $(-1, 0)$ AND THE ORIGIN | F | Q $(4, -1)$ AND R $(4, -4)$ |
| G | M $(,)$ AND N $(3, -5)$ | H | T $(\frac{1}{2}, -\frac{5}{2})$ AND $(-\frac{3}{2}, 1)$ |

2 FIND THE EQUATION OF THE LINE WITH SLOPE m THROUGH THE GIVEN POINT

A $m = \frac{3}{2}$; $P(0, -6)$ **B** $m = 0$; $P\left(\frac{1}{2}, \frac{-}{4}\right)$

C $m = 1\frac{2}{3}$; $P(1, 1)$ **D** $m = -$; $P(0, 0)$

E $m = \sqrt{2}$; $P(\sqrt{2}, -\sqrt{2})$ **F** $m = -1$; $P\left(\frac{1}{3}, \frac{3}{2}\right)$.

3 FIND THE EQUATION OF THE LINE WITH SLOPE m AND Y-INTERCEPT b .

A $m = 0.1$; $b = 0$ **B** $m = -\sqrt{2}$; $b = -1$ **C** $m =$; $b = 2$

D $m = 1\frac{1}{3}$; $b = \frac{-5}{3}$ **E** $m = \frac{-1}{4}$; $b = 5$ **F** $m = \frac{2}{3}$; $b = 1.5$

4 SUPPOSE A LINE HAS X-INTERCEPT a AND Y-INTERCEPT b , $b \neq 0$; SHOW THAT THE EQUATION OF THE LINE IS $\frac{x}{a} + \frac{y}{b} = 1$.

5 FOR EACH OF THE FOLLOWING EQUATIONS, FIND THE SLOPE AND Y-INTERCEPT.

A $\frac{3}{5}x - \frac{4}{5}y + 8 = 0$ **B** $-y + 2 = 0$ **C** $2x - 3y + 5 = 0$

D $x + \frac{1}{2}y - 2 = 0$ **E** $y + 2 = 2(x - 3y + 1)$.

6 A LINE PASSES THROUGH THE POINTS A (5, -1) AND B (-3, 3). FIND:

A THE POINT-SLOPE FORM OF THE EQUATION OF THE LINE.

B THE SLOPE-INTERCEPT FORM OF THE EQUATION OF THE LINE.

C THE TWO-POINT FORM OF THE EQUATION OF THE LINE. WHAT IS ITS GENERAL FORM?

7 FIND THE SLOPE AND Y-INTERCEPT, IF THE EQUATION OF THE LINE IS:

A $\frac{1}{3}x - \frac{2}{3}y + 1 = y + x$ **B** $3(y - 2x) = y + \frac{1}{2}(1 - 2x)$.

8 A TRIANGLE HAS VERTICES AT A (-1, 1), B (1, 3) AND C (3, 1).

A FIND THE EQUATIONS OF THE LINES CONTAINING THE SIDES OF THE TRIANGLE.

B IS THE TRIANGLE A RIGHT-ANGLED TRIANGLE?

C WHAT ARE THE INTERCEPTS OF THE LINE PASSING THROUGH B AND C?

4.4 PARALLEL AND PERPENDICULAR LINES

SLOPES CAN BE USED TO SEE WHETHER TWO NON-VERTICAL LINES IN A PLANE ARE PARALLEL, PERPENDICULAR, OR NEITHER.

FOR INSTANCE, THE LINES $y = x + 3$ AND $y = x$ ARE PARALLEL AND THE LINES $y = x$ AND $y = -x$ ARE PERPENDICULAR. HOW ARE THE SLOPES RELATED?

ACTIVITY 4.8



- 1 WHAT IS MEANT BY TWO LINES BEING PARALLEL? PERPENDICULAR?
- 2 IN FIGURE 4.19, l_1 AND l_2 ARE PARALLEL.
 - A CALCULATE THE SLOPE OF EACH LINE AND THE EQUATION OF EACH LINE.
 - C DISCUSS HOW THEIR SLOPES ARE RELATED.




Figure 4.19




Figure 4.20

- 3 IN FIGURE 4.20 ABOVE, l_1 AND l_2 ARE PERPENDICULAR.
 - A CALCULATE THE SLOPE OF EACH LINE AND THE EQUATION OF EACH LINE.
 - C DISCUSS HOW THEIR SLOPES ARE RELATED.

Theorem 4.1

If two non-vertical lines l_1 and l_2 are parallel to each other, then they have the same slope.

SUPPOSE YOU HAVE TWO NON-VERTICAL LINES WITH SLOPES m_1 AND m_2 AND INCLINATIONS θ_1 AND θ_2 , RESPECTIVELY AS SHOWN IN FIGURE 4.21

IF l_1 IS PARALLEL TO l_2 , THEN $m_1 = m_2$ (WHY?)

CONSEQUENTLY $\tan \theta_1 = \tan \theta_2 = m_2$

State and prove the converse of the above theorem.

What can be stated for two vertical lines? Are they parallel?




Figure 4.21

EXAMPLE 1 SHOW THAT THE LINE PASSING THROUGH A AND B (2, -3) IS PARALLEL TO THE LINE PASSING THROUGH P AND Q (3, 6).

SOLUTION: SLOPE $\overline{AB} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-1)}{2 - (-1)} = \frac{-3 + 1}{2 + 1} = -\frac{2}{3}$

SLOPE $\overline{PQ} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-6 - (-2)}{3 - (-3)} = \frac{-6 + 2}{3 + 3} = -\frac{2}{3}$

SINCE \overline{AB} AND \overline{PQ} HAVE THE SAME SLOPE, PARALLEL TO. $\overline{AB} \parallel \overline{PQ}$

RECALL THAT TWO LINES ARE PERPENDICULAR, IF THEY FORM A RIGHT-ANGLE AT INTERSECTION.

Theorem 4.2

Two non-vertical lines having slopes m_1 and m_2 are perpendicular, if and only if $m_1 \cdot m_2 = -1$.

Proof: SUPPOSE l_1 IS PERPENDICULAR TO l_2

Note: IF ONE OF THE LINES IS A VERTICAL LINE, THEN THE OTHER ONE MUST BE HORIZONTAL WHICH HAS SLOPE ZERO. SO, ASSUME THAT NEITHER LINE IS VERTICAL.

LET m_1 AND m_2 BE THE SLOPES OF l_1 AND l_2 RESPECTIVELY.

LET $R(x_0, y_0)$ BE THE POINT OF INTERSECTION AND CHOOSE $P(x_1, y_1)$ ON l_1 AND $Q(x_2, y_2)$ ON l_2 , RESPECTIVELY.

DRAW TRIANGLES $\triangle QSR$ AND $\triangle RTP$ AS SHOWN IN FIGURE 4.22.

$\triangle QSR$ AND $\triangle RTP$ ARE SIMILAR, (WHY?)

$$\frac{PT}{RT} = \frac{RS}{QS} \quad (\text{WHY?})$$

$$\frac{y_1 - y_0}{x_1 - x_0} = \frac{x_0 - x_2}{y_2 - y_0} = - \left(\frac{x_2 - x_0}{y_2 - y_0} \right)$$

$$\frac{y_1 - y_0}{x_1 - x_0} = \frac{-1}{\frac{y_2 - y_0}{x_2 - x_0}}$$

$$m_1 = - \frac{1}{m_2} \quad \text{OR} \quad m_1 m_2 = -1$$




Figure 4.22

AS AN EXERCISE, START WITH $\frac{QS}{RS} = \frac{RT}{PT}$ AND CONCLUDE THAT $\frac{1}{m_2} = -m_1$

CONVERSELY, YOU COULD SHOW THAT IF TWO LINES l_1 AND l_2 WITH SLOPES m_1 AND m_2 RESPECTIVELY ARE PERPENDICULAR, THEN THE LINES ARE PERPENDICULAR. THIS CAN BE DONE BY REVERSING THE ABOVE PROOF. CONCLUDING THAT THE TWO TRIANGLES ARE SIMILAR. COMPLETE THE PROOF.

EXAMPLE 2 SUPPOSE l_1 PASSES THROUGH P (-1, -3) AND Q (2, 6). FIND THE SLOPE OF

ANY LINE THAT IS:

- A** PARALLEL TO l_1 **B** PERPENDICULAR TO l_1

SOLUTION: THE SLOPE OF l_1 IS

$$m_1 = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - (-3)}{2 - (-1)} = \frac{9}{3} = 3. \text{ SO,}$$

A THE SLOPE OF l_2 PARALLEL TO l_1 IS $m_1 = 3$

B THE SLOPE OF l_2 PERPENDICULAR TO l_1 IS $m_2 = - \frac{1}{m_1} = - \frac{1}{3}$

EXAMPLE 3 FIND THE EQUATION OF THE LINE PASSING THROUGH P(5, 1) AND PERPENDICULAR TO THE LINE $-3y = -7$.

SOLUTION: FROM $-3y = -7$, $y = \frac{1}{3}x + \frac{7}{3}$ SO, $m_1 = \frac{1}{3}$

LET THE SLOPE OF THE REQUIRED LINE BE $m_2 = -1$ GIVES $m_2 = \frac{-1}{m_1} = -3$

THEREFORE THE REQUIRED EQUATION OF THE LINE IS $y = -3x + 14$.

Exercise 4.6

- 1 IN EACH OF THE FOLLOWING, DETERMINE WHETHER THE LINE IS PARALLEL TO OR PERPENDICULAR TO THE LINE THROUGH
 - A A (-1, 3) AND B (2, -2)
P (1, 4) AND Q (-2, 9)
 - B A (-3, 5) AND B (2, -5)
P (-1, 4) AND Q (1, 5).
- 2 FIND THE SLOPE OF THE LINE THAT IS PERPENDICULAR TO l AND Q (-3, -2).
- 3 USE SLOPE TO SHOW THAT THE QUADRILATERAL WITH VERTICES A (-2), B (-3, 1), C (3, 0) AND D (1, -3) IS A PARALLELOGRAM.
- 4 LET l BE THE LINE WITH EQUATION $2x - 3y = 6$. FIND THE SLOPE-INTERCEPT FORM OF THE EQUATION OF THE LINE THAT PASSES THROUGH THE POINT P (2, 1) AND IS
 - A PARALLEL TO l
 - B PERPENDICULAR TO l
- 5 FIND THE EQUATION OF A LINE PASSING THROUGH P AND PERPENDICULAR TO THE LINE l :
 - A $l: 2x - 5y - 4 = 0$; P (-1, 2)
 - B $l: 3x + 6 = 0$; P (4, -6).
- 6 DETERMINE WHICH OF THE FOLLOWING PAIRS OF LINES GIVEN ARE PERPENDICULAR OR PARALLEL OR NEITHER:
 - A $3x - y + 5 = 0$ AND $x + 3y - 1 = 0$
 - B $3x - 4y + 1 = 0$ AND $x - 3y + 1 = 0$
 - C $4x - 10y + 8 = 0$ AND $10x + 6y - 3 = 0$
 - D $2x + 2y = 4$ AND $x + y = 10$.
- 7 FIND THE EQUATION OF THE LINE PASSING THROUGH P AND Q. IS THE LINE
 - A PARALLEL TO THE LINE PASSING THROUGH POINTS A (3, 1) AND B (1, 2)
 - B PERPENDICULAR TO THE LINE JOINING THE POINTS A (4, 2) AND B (1, 2)
 - C PARALLEL TO THE LINE
 - D PERPENDICULAR TO THE LINE.
- 8 DETERMINE THAT THE LINE WITH EQUATION $2x - 3y = 6$ WILL BE:
 - A PARALLEL TO THE LINE WITH EQUATION $4x - 6y = 12$
 - B PERPENDICULAR TO THE LINE WITH EQUATION $2x - 3y = 6$
- 9 SHOW THAT THE PLANE FIGURE WITH VERTICES:
 - A A (6, 1), B (5, 6), C (-4, 3) AND D (-3, -2) IS A PARALLELOGRAM
 - B A (2, 4), B (1, 5), C (-2, 2) AND D (-1, 1) IS A RECTANGLE.
- 10 THE VERTICES OF A TRIANGLE ARE A (1, 8) AND C (6, 4). SHOW THAT THE LINE JOINING THE MID-POINTS OF SIDES AB AND AC IS PARALLEL TO AND ONE-HALF THE LENGTH OF SIDE BC.



Key Terms

analytic geometry	general equation of a line	point-slope form
angle of inclination	horizontal line	slope (gradient)
coordinate geometry	inclination of a line	slope-intercept form
coordinates	mid-point	steepness
equation of a line	non-vertical line	two-point form



Summary

- IF A POINT HAS COORDINATES (x, y) , THEN THE NUMBER x IS CALLED THE **abscissa** OR **horizontal coordinate** AND IS CALLED **the x-coordinate** OF P . THE NUMBER y IS CALLED THE **ordinate** OR **vertical coordinate** OF P .
- THE **distance** d BETWEEN POINTS $P(x_1, y_1)$ AND $Q(x_2, y_2)$ IS GIVEN BY THE FORMULA

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

- THE POINT $R(x_0, y_0)$ DIVIDING THE LINE SEGMENT PQ INTERNALLY, IN THE RATIO $m:n$ IS GIVEN BY

$$R(x_0, y_0) = \left(\frac{nx_1 + mx_2}{n+m}, \frac{ny_1 + my_2}{n+m} \right),$$

WHERE $P(x_1, y_1)$ AND $Q(x_2, y_2)$ ARE THE END-POINTS.

- THE **mid-point** OF A LINE SEGMENT WHOSE END-POINTS ARE $P(x_1, y_1)$ AND $Q(x_2, y_2)$ IS GIVEN BY

$$M(x_0, y_0) = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

- IF $P(x_1, y_1)$ AND $Q(x_2, y_2)$ ARE POINTS ON A LINE, THEN THE **slope (gradient)** OF THE LINE IS GIVEN BY

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

- IF θ IS THE ANGLE BETWEEN THE POSITIVE X-AXIS AND THE LINE PASSING THROUGH THE POINTS $P(x_1, y_1)$ AND $Q(x_2, y_2)$, $x_1 \neq x_2$, THEN THE **slope** OF THE LINE IS GIVEN BY

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \tan \theta$$

- THE GRAPH OF THE EQUATION $x = a$ IS THE **vertical line** THROUGH $(a, 0)$ AND HAS NO SLOPE.
- THE **equation of the line** WITH SLOPE m AND PASSING THROUGH THE POINT $P(x_1, y_1)$ IS GIVEN BY

$$y - y_1 = m(x - x_1)$$

9 THE EQUATION OF THE LINE WITH SLOPE m AND INTERCEPT b IS GIVEN BY

$$y = mx + b$$

10 THE EQUATION OF THE LINE PASSING THROUGH POINTS $P(x_1, y_1)$ AND $Q(x_2, y_2)$ IS GIVEN BY

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1} (x - x_1), x_1 \neq x_2$$

11 THE GRAPH OF EVERY FIRST DEGREE (LINEAR) EQUATION $C = 0$, $A, B \neq 0$ IS A **straight line** AND EVERY STRAIGHT LINE IS A GRAPH OF A FIRST DEGREE EQUATION

12 TWO NON-VERTICAL LINES ARE **parallel** IF AND ONLY IF THEY HAVE THE SAME SLOPE

13 LET ℓ_1 BE A LINE WITH SLOPE m_1 AND ℓ_2 BE A LINE WITH SLOPE m_2 . THEN ℓ_1 AND ℓ_2 ARE **perpendicular** LINES IF AND ONLY IF $m_1 m_2 = -1$.



Review Exercises on Unit 4

1 SHOW THAT THE POINTS $A(-1)$, $B(-1, 1)$ AND $C(\sqrt{3}, \sqrt{3})$ ARE THE VERTICES OF AN EQUILATERAL TRIANGLE

2 FIND THE COORDINATES OF THE THREE POINTS THAT DIVIDE THE LINE SEGMENT $P(-4, 7)$ AND $Q(10, -9)$ INTO FOUR PARTS OF EQUAL LENGTH

3 FIND THE EQUATION OF THE LINE WHICH PASSES THROUGH $P(-4, -2)$ AND $Q(3, 6)$.

4 FIND THE EQUATION OF THE LINE

A WITH SLOPE -3 THAT PASSES THROUGH $P(8, 3)$.

B WITH SLOPE $\frac{1}{2}$ THAT PASSES THROUGH $Q(5)$.

5 IN EACH OF THE FOLLOWING, SHOW THAT THE THREE POINTS ARE VERTICES OF A RIGHT ANGLE TRIANGLE

A $A(0, 0)$, $B(1, 1)$, $C(2, 0)$ **B** $P(3, 1)$, $Q(-3, 4)$, $R(-3, 1)$.

6 FIND THE SLOPE AND INTERCEPT OF THE LINE WITH THE FOLLOWING EQUATIONS:

A $2x - 3y = 4$ **B** $2y - 5x - 2 = 0$

C $5y + 6x - 4 = 0$ **D** $3y = 7x + 1$.

7 FIND THE EQUATION OF THE STRAIGHT LINE PASSING THROUGH P

A PARALLEL TO THE LINE WITH EQUATION $2x$

B PERPENDICULAR TO THE LINE WITH EQUATION $5y$.

8 LET ℓ BE THE LINE THROUGH $A(5)$ AND $B(3, t)$ THAT IS PERPENDICULAR TO THE LINE THROUGH $P(1, 3)$ AND $Q(-4, 2)$. FIND THE VALUE OF t

9 LET ℓ BE THE LINE THROUGH $A(-8)$ AND $B(t, -2)$ THAT IS PARALLEL TO THE LINE THROUGH $P(-2, 4)$ AND $Q(4, -1)$. FIND THE VALUE OF t

10 PROVE THAT THE CONDITION FOR LINES $Ax + C = 0$ AND $ax + by + c = 0$ TO BE PERPENDICULAR MAY BE WRITTEN IN THE FORM

$$Aa + Bb = 0, \text{ WHERE } Bb \neq 0.$$